



BRILL



brill.com/ctoz

The squaretail mullet *Ellochelon vaigiensis* (Quoy & Gaimard, 1825) a complex of cryptic species?

Mohammad Sadegh Alavi-Yeganeh | ORCID: 0000-0002-6246-9504

Tarbiat Modares University, Marine Biology Department, Nur, Mazandaran, Iran

malavi@modares.ac.ir

Javad Ghasemzadeh

Chabahar Maritime University, Fisheries Sciences Department, Chabahar,

Sistan & Balochestan, Iran

Sanaz Kouhi

Tarbiat Modares University, Marine Biology Department, Nur, Mazandaran, Iran

Jean-Dominique Durand | ORCID: 0000-0002-0261-0377

MARBEC, University of Montpellier, IRD, CNRS, cc093, Place E. Bataillon, 34095 Montpellier

Cedex 05, France

RECEIVED 2 NOVEMBER 2022; REVISED AND ACCEPTED 14 MAY 2023; PUBLISHED ONLINE
26 MAY 2023; PUBLISHED IN ISSUE 9 OCTOBER 2023

EDITOR: I. BUJ

Abstract

The teleost family Mugilidae is speciose with uniform morpho-anatomical characteristics, which render species identification difficult. The DNA barcoding technique has, however, proven to be a precise and reliable approach for species delineation. To date, DNA barcoding flags numerous polyphyletic species in Mugilidae that probably correspond to species complexes and that call for further taxonomical investigation. Among these species, the squaretail mullet *Ellochelon vaigiensis* is an interesting case study because, unlike other mullet species, it is easily identified from its unique phenotype. Recent studies of genetic diversity in this monotypic species have revealed two lineages, located either in the Indo-Pacific (Polynesia and Taiwan waters) or along Australian shores. In this study, a third lineage is described in the North of the Indian Ocean, based on nucleotide polymorphisms of Cytochrome *c* oxidase 1 and barcodes available in BOLD and Cytochrome *b*. Despite genetic divergences that exceed the intraspecific threshold, there was no morpho-anatomical difference among specimens of the north Indian Ocean vs. Indo-Pacific

or Australia. These molecular results suggest nominal species of *Ellochelon vaigiensis* within a cryptic species complex.

Keywords

DNA barcoding – integrative taxonomy – morpho-anatomical characters – Mugilidae – phylogeny

Introduction

The Mugilidae family is a taxonomically diverse group of fishes that inhabits all tropical, subtropical, and temperate water bodies of the planet. At a regional scale species diversity is more limited and, based on morphology, eight species of mullet have been reported from the Persian Gulf and Oman Sea (Firouzi et al., 2020). Among these species, the squaretail mullet *Ellochelon vaigiensis* is the only representative of the genus *Ellochelon*, which belongs to the Squalomugilini with two other monotypic genera, namely *Plicomugil* and *Squalomugil* (Xia et al., 2016). It is distinguishable from other mullets by various morphological criteria. These include a broad head, lack of adipose eyelid, truncate caudal fin, pectoral fins without axillary scale that are black in juveniles but have a yellow lower margin in adults, other fins dusky, absence of shelf-like folds inside mouth corners, presence of labial teeth, less than 28 weakly ctenoid and relatively large scales in lateral series, and distinct dark longitudinal strips on lateral scales (Ghasemzadeh, 1998; Harrison & Senou, 1999; Xia et al., 2016).

The species is found in shallow waters (0–5 meters) in lagoons, estuaries, sheltered sandy shorelines and coastal creeks, and will occasionally enter freshwater (McDowall, 1997; Whitfield et al., 2012). Its distribution range encompasses all East African shores north to the Persian Gulf, east to Marshall, Gambier and the Marquesas Islands, north to

southern Japan, south to Western Australia, New South Wales (Australia), New Caledonia, Society Islands, and Rapa the Great Barrier Reef (McDowall, 1997; Harrison & Senou, 1999; Fricke et al., 2020). Over this geographically large distribution, up to ten species have been described and, later, synonymized with *Ellochelon vaigiensis* (Thomson, 1997; Motomura et al., 2010; González-Castro & Ghasemzadeh, 2015). Durand et al. (2012) used mitochondrial sequence polymorphisms, to demonstrate, however, that this species is actually polyphyletic, and revealed two lineages. Considering the degree of divergence (4.8%) between the two lineages, which largely exceeds the accepted level of intraspecific diversity (see Durand et al., 2017), alongside the geographic distribution of the two lineages, Durand and Borsa (2015) suggested conserving the species name for the lineage close to the type locality and provisionally naming the lineage in Australian waters as *Ellochelon* sp. A. In this context, it is important to increase DNA barcoding efforts on squaretail mullet, analyzing more specimens from various localities over its known distribution range in tandem with studies of morpho-anatomy. Using an integrative taxonomy approach, we propose to 1. genetically characterize new specimens from the Persian Gulf and the Oman Sea where the species have only been investigated morphologically (Teimouri & Hesni, 2020; Firouzi et al., 2020); 2. determine their phylogenetic relationships with the existing two lineages, and 3. record

morpho-anatomical characters used to establish the Mugilidae taxonomy for all holotypes and some representative specimens belonging to the different mitochondrial lineages.

Materials and methods

Field sampling

Ten specimens of *Ellochelon vaigiensis* were collected from the Persian Gulf (Bandar Abbas; 27°07'N 56°21'E) and Northern Oman Sea (Gwadar Bay; 25°10'N; 61°29'E; fig. 1) by gillnet and beach seine (fig. 2). The fin clips were fixed in 96% ethanol for molecular studies, and the specimens were transferred to 4% formalin for morphological analysis. Specimens were cataloged in the Aquatic Animal Collection of Tarbiat Modares University (Codes: TAC1066F & TAC1066F for the Persian Gulf specimens and TAC1224F, for the specimens from the Oman Sea.

DNA extraction and sequencing

DNA extraction was carried out by the phenol-chloroform method (Taggart et al., 1992). A Cytochrome *c* oxidase 1 (COI) fragment of 652 base pairs (bp) and a Cytochrome *b* (Cyt-*b*) fragment of 937 bp was amplified using FishF1/FishR1 (Ward et al., 2005) and GluF/ThrR (Machordom & Doadrio, 2001) primers, respectively.

The 40 µl PCR reaction mixes included 20 µl of MyTaq PCR Mastermix (Bioline), 16 µl of

ultrapure water, 0.8 µl of BSA (Euromedex), 0.6 µl of each primer (3 µM), and 2 µl of DNA template. Amplifications were performed in a PTC-100 BIORAD Thermal cycler. The thermal regime for the COI gene fragment consisted of an initial step of 2 min at 92°C followed by 35 cycles of 45 s at 92°C, 45 s at 52°C, and 1 min at 72°C, followed in turn by 5 min at 72°C and then held at 4°C. For the Cyt-*b* gene fragment, the thermal regime consisted of an initial step of 2 min at 94°C followed by 35 cycles of 45 s at 94°C, 1 min at 52°C, and 1 min at 72°C, followed in turn by 7 min at 72°C and then held at 4°C. PCR products were visualized on 1–2% agarose gels and the most intense products were selected for sequencing by MacroGen (<https://dna.macrogen.com>). The quality of sequences (base miscoding) was controlled visually using Chromas software (www.technelysium.com.au/chromas.html).

Morphological analyses

Twenty four morphometric and eight meristic characters (table 1) were measured by caliper to an accuracy of 0.02 mm and counted using a stereomicroscope (Lagler, 1956; Alavi-Yeganeh & Bahmani, 2018), on 10 specimens collected from the Persian Gulf and Oman Sea, and on 24 specimens from the Indo-Pacific: AMS I.22784-035 (52 mm Standard Length; SL), Australia, Cooktown; AMS I.24689-021 (2, 50–52 mm SL), Australia, Darwin, the mouth of Buffalo Creek; AMS 11154 (325 mm SL), Australia, North Queensland; AMS I.9708 (334 mm SL),



FIGURE 1 Squaretail mullet (*Ellochelon vaigiensis*) from Gwadar Bay, Northern Oman Sea, Iran. Total length = 22.3 cm.




FIGURE 2 Sampling locations for squaretail mullet *Ellochelon vaigiensis* from the Persian Gulf (Bandar Abbas) and the Oman Sea (Gwadar Bay).

Australia, Cape York; AMS I.15171 (255 mm SL), Australia, Queensland, Wide Bay; AMS I. 15172 (245 mm SL), Australia, Queensland, Wide Bay; AMS I.20979-004 (132 mm SL), Australia, Queensland, Lizard Island; AMS 1B.4633 (4, 23–33 mm SL), New Guinea, Gum River, Madang; AMS 1B.5523 (95 mm SL), Kiribati, Gilbert Islands; AMS 1B.1514 (192 mm SL), New Guinea, Purdy Archipelago; AMS 1A.1448 (105 mm SL), Australia, Gulf of Carpentaria; AMS: I.34341-004 (40, 24–41 mm SL); Australia, Queensland, Mangrove Channel, Port Clinton; AMS I.20639-002 (3, 119–124 mm SL), Kiribati, Phoenix Islands; AMS I.13755 (2, 58–60 mm SL), Vanuatu, New Hebrides; BMNH 1889.2.1.3764 (182 mm SL), India, Bombay; BMNH 1890.9.23.97 (336 mm SL), Australia, Fraser Island; BMNH 1860.3.19.362 (243 mm SL), Malaysia, Penang; BMNH 1872.4.6.13 (365 mm SL), Tahiti; AMS I.34341-004 (3, 30–38 mm SL), Australia, Queensland, Mangrove Channel, Port Clinton; AMS I.22784-035 (52 mm SL), Australia, Cooktown; MQU F.1040 (4, 77–83 mm SL), Australia, Queensland, Lwize Creek, Hector; NTM S.10811-004 (76 mm SL), Australia, Dampier Archipelago; QM I.7461 (92 mm SL), Australia, Lindeman Island; WAM

P.29916-008 (3, 61–70 mm SL). A multivariate analysis (Principal Component Analysis; PCA) was carried out by PAST Ver. 2.17 to identify the most effective character to differentiate morphological differences between two groups of specimens.

Phylogenetic analyses

The dataset used for the phylogenetic investigation consisted of three COI and three Cyt-b sequences from this study which were uploaded to GenBank (MT843281-3 and MT882672-4, respectively), with all sequences stored in the BOLD DNA barcodes library (<http://boldsystems.org>, consultation on Nov. 2021) associated with the name "*Ellochelon vaigiensis*" and belonging to three Barcode Index Numbers (BINS) BOLD: AAC9398, BOLD: AAU0553, BOLD: ACK7668 and five available congener Cyt-b sequence from the GenBank were extracted (table 2). A fourth BIN (BOLD: ABY5947) is associated with the name "*Ellochelon vaigiensis*" but was not considered in our final dataset as only one member of the 20 members is identified as *Ellochelon vaigiensis*, and certainly corresponds to a misidentification considering

TABLE 1 Comparison of morphometric and meristic characters for specimens of *Ellocheilon vaigiensis* from Indo-Pacific and Northern Indian Ocean.

Character	Indo-Pacific (N = 20)	Northern Indian Ocean (N = 7)
Morphometric characters		
Total length	96.5–131.5 (107.1 ± 9.8)	70.2–220.0 (106.8 ± 45.1)
Standard length	76.5–113.4 (84.9 ± 10.6)	57.0–77.0 (85.5 ± 36.4)
Head length/ SL %	26.2–31.4 (30.5 ± 1.6)	26.3–32.8 (29.2 ± 2.0)
Pectoral fin length/ SL %	22.8–26.0 (24.9 ± 1.0)	22.1–24.9 (23.0 ± 0.8)
Body depth (Max)/ SL %	27.4–30.3 (29.0 ± 0.8)	24.0–28.8 (27.3 ± 1.5)
Body depth (Min)/ SL %	13.6–15.3 (14.5 ± 0.5)	12.6–15.0 (14.3 ± 0.7)
Pre dorsal (1) length/ SL %	47.8–56.6 (54.3 ± 2.4)	51.9–54.9 (53.5 ± 1.0)
Dorsal (1) to caudal base distance/ SL %	42.8–48.3 (46.9 ± 1.6)	45.7–50.0 (48.1 ± 1.3)
Pre dorsal (2) length/ SL %	70.3–80.7 (77.8 ± 2.9)	75.4–82.3 (78.6 ± 1.8)
Pre pelvic length/ SL %	36.0–44.7 (42.2 ± 2.4)	37.7–43.0 (39.8 ± 1.6)
Pre anal length/ SL %	66.1–78.0 (74.7 ± 3.2)	69.7–75.8 (73.1 ± 1.8)
Caudal peduncle length/ SL %	11.4–14.1 (12.7 ± 0.9)	12.3–15.5 (13.7 ± 1.2)
Snout length/Head length%	21.2–30.0 (26.0 ± 3.1)	16.0–24.5 (20.6 ± 3.0)
Eye length/Head length %	25.0–27.4 (26.0 ± 0.9)	20.4–25.9 (23.8 ± 1.6)
Interorbital length/Head length%	47.5–53.2 (49.1 ± 1.8)	42.8–54.2 (49.0 ± 3.6)
Postorbital length/Head length%	47.0–52.5 (48.9 ± 1.7)	42.8–58.5 (53.0 ± 4.9)
Mouth width/Head length%	38.5–46.1 (40.1 ± 2.2)	32.1–43.5 (36.7 ± 4.4)
Body width/Head length%	71.5–81.1 (74.9 ± 2.8)	65.8–82.8 (74.9 ± 5.4)
Caudal ped width/Head length%	14.1–20.9 (17.2 ± 2.3)	11.2–18.3 (13.4 ± 1.9)
Dorsal fin height/Head length%	48.1–57.2 (53.3 ± 3.0)	42.9–54.2 (47.6 ± 4.0)
Dorsal fin 2 height/Head length%	60.2–71.7 (64.6 ± 3.6)	63.7–73.1 (66.7 ± 3.4)

TABLE 1 Comparison of morphometric and meristic characters for specimens of *Ellochelone vaigiensis* from Indo-Pacific and Northern Indian Ocean. (cont.)

Character	Indo-Pacific (N = 20)	Northern Indian Ocean (N = 7)
Anal fin height/Head length%	61.7–5.1 (69.3 ± 4.0)	66.0–79.6 (71.6 ± 4.0)
Pectoral fin length/Head length%	76.8–6.9 (81.9 ± 2.7)	69.5–94.6 (79.1 ± 7.7)
Ventral fin length/Head length%	60.6–71.7 (64.4 ± 2.9)	58.0–71.4 (62.8 ± 4.4)
Meristic characters		
Midlateral scales	26–27 (26.8 ± 0.4)	26–27 (26.4 ± 0.5)
Vertebral count	24 (24 ± 0)	24 (24 ± 0)
Second dorsal fin rays	8 (8 ± 0)	7–8 (7.8 ± 0.4)
Anal fin spine	3 (3 ± 0)	3 (3 ± 0)
Anal fin rays	8 (8 ± 0)	8–9 (8.4 ± 0.5)
Pectoral fin rays	15–16 (15.6 ± 0.5)	15–16 (15.3 ± 0.5)
Pyloric caeca number	13–24 (18.0 ± 3.5)	12–15 (13.3 ± 1.3)
Gill rakers	43–47 (43.9 ± 1.4)	47777777–51 (48.5 ± 1.7)

TABLE 2 Voucher specimens and accession numbers used in the present molecular analyses for COI & Cyt-b gene sequences.

Accession number	Voucher number	Bold	BIN	Location
COI				
JQ060443	CSIRO H4307-01	ANGBF5311-12	BOLD:ACK7668	Australia, Unspecified location
n.a.	CSIRO H4307-01	FOAC173-05	BOLD:ACK7668	Australia, Queensland
n.a.	n.a.	FOAC174-05	BOLD:ACK7668	Australia, Queensland
KP194416	n.a.	LIFS409-08	BOLD:ACK7668	Australia, Queensland, Lizard Island
KP194221	n.a.	LIFS410-08	BOLD:ACK7668	Australia, Queensland, Lizard Island
KP194088	CSIRO H 7589-02	LIFS411-08	BOLD:ACK7668	Australia
KP194965	n.a.	LIFS412-08	BOLD:ACK7668	Australia, Queensland, Lizard Island
KP194097	n.a.	LIFS696-08	BOLD:ACK7668	Australia, Queensland, Lizard Island
LC114151	n.a.	ANGBF37664-19	BOLD:AAU0553	Iran
MT843281	TAC1224F-01	Pending	BOLD:AAU0553	Iran/ Oman (This study)
MT843282	TAC1224F-02	Pending	BOLD:AAU0553	Iran/ Oman (This study)
MT843283	TAC1066F	Pending	BOLD:AAU0553	Iran Persian Gulf (This study)
KY849517	n.a.	ANGBF37663-19	BOLD:AAU0553	Peninsular Malaysia
n.a.	n.a.	FOAN718-11	BOLD:AAU0553	Indonesia, Nusa Tenggara Barat, Lombok
MT885104	MZB-BIF00353	BIFB328-13	BOLD:AAU0553	Indonesia, Jawa Barat, Kabupaten Sukabumi
KU943215	ASIZP0805165	ZOSKT368-16	BOLD:AAC9398	Taiwan, Dongsha atoll
KU943216	ASIZP0805166	ZOSKT369-16	BOLD:AAC9398	Taiwan, Dongsha atoll
MT885092	MZB-BIF5603	BIFD4722-16	BOLD:AAC9398	Indonesia, Maluku, Ambon Island
MT885017	MZB-BIF5604	BIFD4723-16	BOLD:AAC9398	Indonesia, Maluku, Ambon Island
MT885071	MZB-BIF5593	BIFZI514-17	BOLD:AAC9398	Indonesia, Maluku, Ambon Island
MT885054	MZB-BIF5601	BIFZI522-17	BOLD:AAC9398	Indonesia, Maluku, Ambon Island
MT885000	MZB-BIF5606	BIFZI527-17	BOLD:AAC9398	Indonesia, Maluku, Ambon Island

TABLE 2 Voucher specimens and accession numbers used in the present molecular analyses for COI & Cyt-b gene sequences. (cont.)

Accession number	Voucher number	Bold	BIN	Location
MT885155	MZB-BIF5610	BIFZI531-17	BOLD: AAC9398	Indonesia, Maluku, Ambon Island
MT884991	MZB-BIF5612	BIFZI533-17	BOLD: AAC9398	Indonesia, Maluku, Ambon Island
MT884960	MZB-BIF5616	BIFZI537-17	BOLD: AAC9398	Indonesia, Maluku, Ambon Island
MT885113	MZB-BIF5617	BIFZI538-17	BOLD: AAC9398	Indonesia, Maluku, Ambon Island
MT884990	n.a.	BIFV026-19	BOLD: AAC9398	Indonesia, Bali
MT885051	MZB-BIF6536	BIFZI665-17	BOLD: AAC9398	Indonesia, Riau, Pekanbaru
JQ060444	n.a.	ANGBF5186-12	BOLD: AAC9398	Indonesia, West Papua, Batanta Island
MT885114	RCO_REF-17-0839	WPRFM001-19	BOLD: AAC9398	Indonesia, West Papua, Kaimana
MT884955	RCO_REF-17-0840	WPRFM002-19	BOLD: AAC9398	Indonesia, West Papua, Kaimana
MT885027	RCO_REF-17-0841	WPRFM003-19	BOLD: AAC9398	Indonesia, West Papua, Kaimana
MT884970	RCO_REF-17-0842	WPRFM004-19	BOLD: AAC9398	Indonesia, West Papua, Kaimana
MT885001	RCO_REF-17-1456	WPRFM005-19	BOLD: AAC9398	Indonesia, West Papua, Kaimana
MT885049	RCO_REF-17-1457	WPRFM006-19	BOLD: AAC9398	Indonesia, West Papua, Kaimana
KP194065	n.a.	LIFS695-08	BOLD: AAC9398	Australia, Queensland, Lizard Island
JQ431702	MNHN_2008-688	MBFA489-07	BOLD: AAC9398	French Polynesia, Society Islands
Q431701	MNHN_2008-689	MBFA490-07	BOLD: AAC9398	French Polynesia, Society Islands
MK657139	USNM:FISH:401336	GAMBA371-12	BOLD: AAC9398	French Polynesia, Tuamotu-Gambier
MK657182	USNM:FISH:401333	GAMBA372-12	BOLD: AAC9398	French Polynesia, Tuamotu-Gambier
MK657095	USNM:FISH:401135	GAMBA373-12	BOLD: AAC9398	French Polynesia, Tuamotu-Gambier
Q060445	n.a.	ANGBF5310-12	BOLD: AAC9398	French Polynesia, Tahiti
Cyt-b				
Accession number	Voucher number	Location		
KF375124	n.a.	Indonesia		
KF375125	n.a.	China		

TABLE 2 Voucher specimens and accession numbers used in the present molecular analyses for COI & Cyt-b gene sequences. (cont.)

Accession number	Voucher number	Location
JQ060187	n.a.	Batanta Island, West Papua
JQ060188	n.a.	Tahiti, French Polynesia
JQ060186	CSIRO-H4307-01	Unspecified location, Australia
MT882672	TAC1224F-01	Iran/Oman (This study)
MT882673	TAC1224F-02	Iran/Oman (This study)
MT882674	TAC1066F	Iran Persian Gulf (This study)

member's pictures associated to this BIN. Alignment was executed by Muscle in MEGA 7 (Kumar et al., 2016). Also, the genetic distance was estimated with the Kimura-2-parameter (K2P) model by MEGA 7 (Kumar et al., 2016). The best substitution model for our datasets was estimated using the method implemented in jModeltest Ver. 2.1.10 (Darriba et al., 2012) and considering the Akaike information criterion (AIC) score. The phylogenetic tree was constructed using maximum likelihood (raxmlGUI; Edler et al., 2021) and Bayesian likelihood (Mr. Bayes; Ronquist & Huelsenbeck, 2003) methods. The Hasegawa, Kishino and Yano with Gamma distribution (HKY + G) and generalized time-reversible with Gamma distribution (GTR + G) substitution models were used for the construction of COI and Cyt-b trees, respectively.

Initial trees for the heuristic search were obtained automatically by applying Neighbor-Joining and BioNJ algorithms to a matrix of pairwise distances estimated using the Maximum Composite Likelihood (MCL) approach and then selecting the topology with a superior log likelihood value. Codon positions included were 1st+2nd+3rd. All positions containing missing data were eliminated. A bootstrap test with 500 replications was performed for the ML (Maximum Likelihood) tree and two simultaneous runs with four MCMC chains were used for 10^7 generations in BL (Bayesian Likelihood) analysis. Subsampling trees and parameters were saved every 1000 generations. A sequence of *Plicomugil labiosus* (JQ060620) was used as an outgroup due to its phylogenetic proximity with *Ellochelon vaigiensis* (Durand et al., 2012) for the COI tree. Also, for the Cyt-b tree, a sequence from *Plicomugil labiosus* (KF375164) was used as an outgroup. Three DNA sequences-based methods of species delimitation including Automatic Barcode Gap Discovery (ABGD) (<https://bioinfo.mnhn.fr/abi/public>)

/abgd/abgdweb.html) (Puillandre et al., 2012), Poisson tree processes (PTP; executed on the PTP web server; <http://species.h-its.org>) (Zhang et al., 2013) and Refined Single Linkage (RESL) (Ratnasingham & Hebert, 2013), were used as species delimitation tools for COI dataset.

Results

None of the morphometric and meristic characters was diagnostic and all had overlapped ranges (table 1). Despite these overlaps in ranges, multivariate analysis (PCA) revealed that pectoral fin length, first dorsal fin height, snout length, postorbital length, anal fin length, body width and a meristic character (pyloric caeca number) were the most effective morphometric characters to discriminate between specimens from Indo-Pacific and Northern Indian Ocean (fig. 3, table 3).

Alignment of COI gene fragments over 551 bp for 42 specimens identified as *Ellochelon vaigiensis* revealed 49 variables and 45 parsimony-informative substitutions. The phylogenetic tree highlighted three lineages in agreement with BIN delineated by the RESL algorithm (Ratnasingham & Hebert, 2013). Three clades supported strongly with bootstrap test (>95) and posterior probabilities values (>0.99). All specimens collected

in the Persian Gulf and the Oman Sea belong to the BIN BOLD: AAU0553 (fig. 4). The two other BINs included specimens used in the phylogenetic investigation of Durand et al. (2012). The genetic distance between *E. vaigiensis* BIN ranges between 5.5% to 6.1%. IntraBIN mean divergences ranged between 0.1% to 0.2% (K2P). None of the haplotypes belonging to these BIN were separated by more than two mutations, with the exception of one haplotype observed in a single specimen from Indonesia (BIFZI665-17) of the BIN BOLD: AAC9398 presents seven mutations to the second haplotype shared by all specimens of this lineage. Using ABGD or PTP, this divergent haplotype is considered a Molecular Operational Taxonomic Unit (MOTU). This difference is the only difference between RESL and the two other species delimitations models (fig. 4). The geographic distribution of the three BINs is only partially overlapping. The BIN BOLD: AAC9398 is present in a specimen collected in the Pacific from Taiwan to Lizard Island (Australia) and from Indonesia to French Polynesia. The BIN BOLD: ACK7668 is present in a specimen collected in Australia, while the BIN BOLD: AAU0553 is observed in the Indian Ocean from the Persian Gulf to Indonesia (Lombok) (fig. 4). Similarly, a phylogenetic tree based on Cyt-b sequence (937 bp) revealed three distinct lineages from southern waters of Iran (lineage I), Australia




FIGURE 3 A scatterplot for PCA based on morphometric and meristic characters of two *Ellochelon vaigiensis* groups from the northern Indian Ocean (square) and Indo-Pacific (circle) areas.

TABLE 3 Loading values of two main Principal components of multivariate analysis (PCA) for morphometric and meristic data of squaretail mullets from Northern Indian Ocean (n = 10) and Indo-Pacific area (n = 24). Characters with higher loading values are presented with bold font.

Character	Loadings	
	Component 1	Component 2
Morphometrics % SL		
Head length	0.123	0.061
Pectoral fin length	0.091	-0.062
Body depth (Max)	0.128	0.010
Body depth (Min)	-0.001	-0.009
Pre dorsal (1) length	0.076	0.003
Dorsal (1) to caudal base distance	-0.063	-0.005
Pre dorsal (2) length	-0.047	-0.164
Pre pelvic length	0.164	-0.072
Pre anal Length	0.132	-0.031
Caudal peduncle length	-0.054	-0.027
Morphometrics % HL		
Snout length	0.325	-0.033
Eye length	0.080	-0.155
Interorbital length	0.036	0.116
Postorbital length	-0.202	0.414
Mouth width	0.236	-0.290
Body width	0.080	0.337
Caudal ped width	0.209	-0.099
First dorsal fin height	0.455	0.249
Second dorsal fin height	-0.222	0.125
Anal fin height	-0.240	0.321
Pectoral fin length	0.362	0.572
Ventral fin length	0.074	0.056
Meristic		
Midlateral scales	0.012	-0.029
Vertebral count	0.001	0.000
Second dorsal fin rays	0.010	-0.001
Anal fin spine	0.000	0.000
Anal fin rays	-0.025	-0.002
Pectoral fin rays	0.023	0.006
Pyloric caeca number	0.345	-0.020
Gill rakers	-0.263	0.169
% of variance	24.7	16.4




FIGURE 4 Phylogenetic diversity of *Ellochelone vaigiensis* based on COI gene sequence data. In (A), the phylogenetic relationships were inferred by using the BL & ML method. The tree is drawn to scale, with branch lengths measured in the number of substitutions per site. Bootstrap percentage values for ML and posterior probability for BL analysis are indicated top and under the nodes, respectively. Haplotypes grouping by ABGD, PTP, and RESL delimitation models are presented with different colors. In (B), a distribution map of locations for three main clade haplotypes is provided. Haplotypes in BOLD format are reference sequences published in Durand and Borsa (2015).

(lineage II) & Polynesia + West Papua + China + Indonesia (lineage III). Clades supported with bootstrap test (>99) and posterior probabilities values (>1.0) (fig. 5).

Discussion

In this study, haplotypes of the nominal species *Ellochelone vaigiensis* were structured into

three independent evolutionary lineages, revealing a novel third lineage in the Indian Ocean. Among the lineages, two were already present in the Mugilidae phylogeny produced by Durand et al. (2012) using an extremely limited sample size; 3 individuals were collected in Australia, Indonesia, and French Polynesia. Recently, a COI haplotype from specimens morphologically identified as *Ellochelone vaigiensis* was reported from the Pakistan coast,




FIGURE 5 The phylogenetic relationship among identified Cyt-b haplotypes of *Ellochelon vaigiensis*. The phylogenetic tree is constructed based on BL & ML methods based on GTR + G model. The numbers at nodes are related to Bayesian posterior probabilities (up) and bootstrap support (down).

which was closely related to specimens assigned by BOLD to the BIN BOLD: AAU0553 (BIFB328-13 from Indonesia) (Hasan et al., 2021) and concordant with the result of this study that identified more specimens belonging to this BIN (fig. 4).

Geographic distributions of the three lineages have been established using additional samples from the Oman Sea and the Persian Gulf, plus DNA barcodes available in the BOLD system. This has revealed a clear phylogeographic pattern, with the three lineages that rapidly diverged from a common ancestor with an apparently parapatric distribution. The suture zone among these different lineages is located in the South of the coral triangle (in Indonesia and North Australia) which suggests a Sunda Shelf (the Indo-Malayan region) origin. This part of the world is known to have a complex geologic history and harbors a hot-spot of marine biodiversity (Tornabene et al., 2015). During the Pleistocene era, this area experienced repeated large-scale eustatic sea-level movements that directly impacted shorelines and coral reefs, and thus the biological connectivity of marine organisms inhabiting it (Voris, 2000; Carpenter et al., 2011; Ludt & Rocha, 2015). Biogeographic barriers created

by sea-level lowering as well as the drastic reduction of habitable shelf area would have prevented gene flow and favored genetic population bottlenecks in *Ellochelon vaigiensis*. This process would explain the low intra-lineage genetic diversity as well as the strong lineage sorting. This phylogeographic signature is frequently observed in marine Indo-Pacific teleost species such as *Dascyllus trimaculatus*, *Decapterus russelli*, *Epinephelus areolatus*, *Epinephelus fasciatus* (for a review, Carpenter et al., 2011; Ludt & Rocha, 2015; Durand et al., 2020), which mirror the magnitude of these geological events in their genetic architecture. Curiously, in the case of *Ellochelon vaigiensis* this process led to three lineages, suggesting that sea-level lowering probably created barriers that led to, at least, three refugia. It remains to be investigated whether the genetic isolation was long enough, or evolutionary forces strong enough, to achieve full speciation in *Ellochelon vaigiensis*.

Considering the level of divergence observed among these lineages, close to or higher than 6% (using the partial COI gene sequence variation), Durand and Borsa (2015) questioned the taxonomic status of this species as well as other polyphyletic

mullet species. The presence of deep genetic divergence between COI haplotypes within a nominal fish species may flag overlooked species (Zemlak et al., 2009) and barcoding gap delineation has been proposed as a method for preliminary identification (Puillandre et al., 2012; Ratnasingham & Hebert, 2013). In their review of fish DNA barcodes generated during the FISH-BOL program, Zemlak et al. (2009) noted that 93% of intraspecific divergences were <1%. Specifically, for the Mugilidae family, Durand et al. (2017) also noted that the distribution of pairwise nucleotide distances based on COI barcodes exhibited a first peak <1% that corresponded to the nucleotidic diversity observed inside MOTUS recovered with the ABGD method (Puillandre et al., 2012). This observation led Durand and Borsa (2015) to propose an interim taxonomic nomenclature for Mugilidae where each genetic lineage (MOTU) is acknowledged and considered as a candidate species. Implicitly, this interim taxonomic nomenclature based on DNA barcode gaps called for further morpho-anatomical investigation, since genetic characters are useless for fish applied taxonomy that relies on morphological and meristic differences. Despite this conclusion, few taxonomic investigations of Mugilid diversity have adopted an integrative approach. A review of *Mugil* species inhabiting South America highlighted that, in *Mugil* sp. N *sensu* Durand and Borsa (2015), there was a tenuous series of meristic characteristics justifying the creation of the species *Mugil margaritae* by Menezes et al. (2015). More recently, Thieme et al. (2022) demonstrated, both genetically and morphologically, that *Chelon* sp. A, which was described in South Africa (Durand & Borsa, 2015), was in fact *Chelon persicus* as described originally in the Persian Gulf (Senou et al., 1996).

Specimens collected from the northern Indian Ocean did not show any diagnostic

morphometric differences from specimens from the Indo-Pacific, although multivariate analysis highlighted two groups of individuals consistent with the genetic delimitation. Similarly, Annisa et al. (2021) compared two populations of *E. vaigiensis* from southern Indonesia (Arafura Sea) and a population from northern Indonesia (Pacific Ocean) morphologically. They did not find any significant morphological differences but multivariate analysis suggested the possible presence of cryptic species in *E. vaigiensis*. That study also emphasized the need for molecular approaches to resolve the taxonomic status.

Integrative taxonomy is not, however, a guarantee of species concept reconciliation (morphology vs. genetics), as demonstrated in the present study where no morphometric or meristic differentiation was highlighted among *Ellochelon vaigiensis* lineages. Morphological conservatism in the Mugilidae has frequently been highlighted and explains why the nomenclature of the family is still largely debated, even at the genus level (for reviews, see Durand, 2016; Xia et al., 2016). In such a situation, it is not surprising to find cryptic species in nominal species such as *Ellochelon vaigiensis*. While that does not help applied taxonomy, an interim taxonomy is still necessary because reliable and relevant for conservation or evolutionary investigations must consider all biodiversity components, especially when a strong biogeographic pattern is identified. For the sake of standardization, the genus name + the BIN should be used to name lineages (OTU) in a species complex; the species name being conserved for the lineage observed at the type locality and if only one lineage is present. In the present situation, the name *Ellochelon vaigiensis* may be maintained only for specimens belonging to the BIN BOLD: AAC9398, while the two other species may be named *Ellochelon* [BOLD: ACK7668] and *Ellochelon* [BOLD: AAU0553].

for the species present in Australia and the Indian Ocean, respectively.

Acknowledgments

We acknowledge H. Zadabbas, M. Khajavi & S. Maric for helping with sampling and David McKenzie for English edition. In addition, we are grateful to two anonymous reviewers who provided valuable comments. We are thankful to Tarbiat Modares University, Chabahar Maritime University and IRD for financial support. The research was approved by the Ethics Committee of Tarbiat Modares University. Data are available only on personal request to the authors. The studied fishes are deposited in Tarbiat Modares Aquatic Animal collection, Tarbiat Modares University, Nur, (TAC1066F & TAC1224F).

References

- Alavi-Yeganeh, M.S. & Bahmani, G. (2018) First Record of Goldstripe Ponyfish, *Karalla daura* (Cuvier, 1829) (Perciformes: Leiognathidae) from the Iranian Coast of the Oman Sea. *Acta Zool. Bulgar.*, 70, 121–124.
- Annisa, A., Omar, S.B.A., Marasabessy, F., Ernawati, E., Kagiling, R., Manangkalangi, E., Tuhumury, R.A.N., Mandala, W.F., Ohee, H.L., Melmambessy, E.H.P., Mote, N., Hocdé, R., Pouyau, L., & Kadarusman, K. (2022). Taxonomic Note on *Ellochelon vaigiensis* (Quoy & Gaimard, 1825) (Mugilidae: Mugiliformes) from West New Guinea, Indonesia. *Trends Sci.*, 19(17), 5750–5750.
- Carpenter, K.E., Barber, P.H., Crandall, E.D., Ablan-Lagman, M.C.A., Ambariyanto, A., Mahardika, G.N., Manjaji-Matsumoto, B.M., Juinio-Meñez, M.A., Santos, M.D., Starger, C.J., & Toha, A.H.A. (2011) Comparative phylogeography of the coral triangle and implications for marine management. *J. Mar. Biol.*, 2011, 1–14. <https://doi.org/10.1155/2011/396982>.
- Darriba, D., Taboada, G. L., Doallo, R., & Posada, D. (2012). jModelTest 2: more models, new heuristics and parallel computing. *Nat. Methods.*, 9(8), 772–772.
- Durand, J.D. (2016) Implications of Molecular Phylogeny for the Taxonomy of Mugilidae. In: D. Crosetti D. and Blaber S. (Eds.) *Biology, Ecology and Culture of Grey Mulletts (Mugilidae)* Taylor and Francis Group, Boca Raton, pp 22–41.
- Durand, J.D. & Borsa, P. (2015) Mitochondrial phylogeny of grey mullets (Acanthopterygii: Mugilidae) suggests high proportion of cryptic species. *J. Crust. Biol.*, 338, 266–277.
- Durand, J.D., Shen, K.N., Chen, W.J., Jamandre, B.W., Blel, H., Diop, K., Nirchio, M., Garcia de Leon, F.J., Whitfield, A.K., Chang, C.W. & Borsa, P. (2012) Systematics of the grey mullet (Teleostei: Mugiliformes: Mugilidae): molecular phylogenetic evidence challenges two centuries of morphology-based taxonomic studies. *Mol. Phylogenet. Evol.*, 64, 73–92. <https://doi.org/10.1016/j.ympev.2012.03.006>.
- Durand, J.D., Hubert, N., Shen, K.N. & Borsa, P. (2017) DNA barcoding grey mullets. *Rev. Fish Biol. Fisher.*, 27, 233–243. <https://doi.org/10.1007/s1160-016-9457-7>.
- Durand, J.D., Pham, M.H., Tran, T.T.V., Hoang, D.H. & Van Vo, Q. (2020) Sorting the wheat from the chaff: a review of BINs associated with groupers of Vietnam and the implications for species identification from DNA barcoding. *Mar. Biodivers.*, 50, 1–23. <https://doi.org/10.1007/s12526-020-01051-1>.
- Edler, D., Klein, J., Antonelli, A., & Silvestro, D. (2021). raxmlGUI 2.0: a graphical interface and toolkit for phylogenetic analyses using RAxML. *Methods Ecol. Evol.*, 12, 373–377.
- Firouzi, N., Ghasemzadeh, J. & Alavi-Yeganeh, M.S. (2020) Morphological characters of scales in ten Mullet species from Iran. *Appl. Biol.*, 33, 111–121. (English Abstract).

- Fricke, R., Eschmeyer, W.N. & Van der Laan, R. (2020) Eschmeyers Catalog of Fishes: Genera, Species, References (<http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>). Electronic version Accessed 2 November 2021.
- Ghasemzadeh, J. (1998) Phylogeny and systematics of Indo-Pacific mullets (Teleostei: Mugilidae) with special reference to the mullets of Australia. Dissertation, Macquarie University.
- González-Castro, M. & Ghasemzadeh, J. (2015) Morphology and morphometry based taxonomy of Mugilidae. In: Crosetti D. & Blaber S. (Eds.) Biology, ecology and culture of grey mullet (Mugilidae). Pp. 1–21. CRC Press Taylor & Francis Group, Boca Raton, USA.
- Hammer, Ø., David, A.T.H. & Ryan, D.P. (2001) PAST: Paleontological statistics software package for education and data analysis. *Palaeontol. Electron.*, 4, 1–9.
- Harrison, I.J. & Senou, H. (1999) Order Mugiliformes. Mugilidae. Mulletts. In: Carpenter K.E. and Niem V.H. (Eds.) FAO species identification guide for fishery purposes. The living marine resources of the Western Central Pacific. Volume 4. Bony fishes' part 2 (Mugilidae to Carangidae). pp. 2069–2108. FAO, Rome.
- Hasan, A., Siddiqui, P.J.A., Amir, S.A. & Durand, J.D. (2021) DNA Barcoding of Mulletts (Family Mugilidae) from Pakistan Reveals Surprisingly High Number of Unknown Candidate Species. *Diversity*, 13, 232. <https://doi.org/10.3390/d13060232>.
- Kumar, S., Stecher, G. & Tamura, K. (2016) MEGA 7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol. Phylogenet. Evol.*, 33, 1870–1874. <https://doi.org/10.1093/molbev/msw054>.
- Lagler, K.F. (1956) Freshwater fishery biology. W.M.C. McGraw Hill, 2nd Edition.
- Ludt, W.B. & Rocha, L.A. (2015) Shifting seas: The impacts of Pleistocene sea-level fluctuations on the evolution of tropical marine taxa. *J. Biogeogr.*, 42, 25–38. <https://doi.org/10.1111/jbi.12416>.
- Machordom, A., & Doadrio, I. (2001) Evidence of a Cenozoic Betic–Kabalian connection based on freshwater fish phylogeography (*Luciobarbus*, Cyprinidae). *Mol. Phylogenet. Evol.*, 18, 252–263.
- McDowall, R.M. (1997) The evolution of diadromy in fishes (revisited) and its place in phylogenetic analysis. *Rev. Fish Biol. Fisher.*, 7, 443–462. <https://doi.org/10.1023/A:1018404331601>.
- Menezes, N.A., Nirchio, M., Oliveira, C.D. & Siccharamirez, R. (2015) Taxonomic review of the species of *Mugil* (Teleostei: Perciformes: Mugilidae) from the Atlantic South Caribbean and South America, with integration of morphological, cytogenetic and molecular data. *Zootaxa*, 3918, 1–38. <https://doi.org/10.11646/zootaxa.3918.1.1>.
- Motomura, H., Kuriwa, K., Katayama, E., Senou, H., Ogihara, G., Meguro, M., Matsunuma, M., Takata, Y., Yoshida, T., Yamashita, M., Kimura, S., Endo, H., Murase, A., Iwatsuki, Y., Sakurai, Y., Harazaki, S., Hidaka, K., Izumi, H. & Matsuura, K. (2010) Annotated checklist of marine and estuarine fishes of Yaku-Shima Island, Kagoshima, southern Japan. In: Motomura H, Matsuura K (Eds) Fishes of Yaku-Shima Island. pp. 65–247. National Museum of Nature and Science, Tokyo.
- Puillandre, N., Lambert, A., Brouillet, S. & Achaz, G. (2012) ABGD, Automatic Barcode Gap Discovery for primary species delimitation. *Mol. Ecol.*, 21, 1864–1877. <https://doi.org/10.1111/j.1365-294X.2011.05239.x>.
- Ratnasingham, S. & Hebert, P.D.N. (2013) A DNA-based registry for all animal species: the barcode index number (BIN) system. *PLoS One*, 8, e66213. <https://doi.org/10.1371/journal.pone.0066213>.
- Ronquist, F. & Huelsenbeck, J.P. (2003) MRBAYES 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics*, 19, 1572–1574.

- Senou, H., Randall, J.E. & Okiyama, M. (1996) *Chelon persicus*, a new species of mullet (Perciformes, Mugilidae) from the Persian Gulf. *Bull. Kanagawa Prefect. Mus. Nat. Sci.*, 25, 71–78.
- Taggart, J.B., Hynes, R.A., Prodöuhl, P.A. & Ferguson, A. (1992) A simplified protocol for routine total DNA isolation from salmonid fishes. *J. Fish Biol.*, 40, 963–965. <https://doi.org/10.1111/j.1095-8649.1992.tb02641.x>.
- Teimori, A. & Hesni, M.A. (2020) Morphological Characteristics of Squaretail Mullet *Ellochelon vaigiensis* (Quoy & Gaimard 1825), a Rare *Mugil* Species Collected from the Iranian Waters of the Persian Gulf (Teleostei: Mugiliformes). *Thalassas*, 36, 405–413. <https://doi.org/10.1007/s41208-020-00217-9>.
- Thieme, P., Bogorodsky, S.V., Alpermann, T.J., Whitfield, A.K., Freitas, R. & Durand, J.D. (2022) Contributions to the taxonomy of the mugilid genus *Chelon* Artedi (Teleostei: Mugilidae), with a major review of the status of *C. persicus* Senou, Randall & Okiyama, 1995. *Zootaxa*, 5188. <https://doi.org/10.11646/zootaxa.5188.1.1.1.42>.
- Thomson, J.M. (1997) The Mugilidae of the world. *Mem. Queensl. Mus.*, 41, 457–562.
- Tornabene, L., Valdez, S., Erdmann, M. & Pezold, F. (2015) Support for a 'Center of Origin' in the Coral Triangle: Cryptic diversity, recent speciation, and local endemism in a diverse lineage of reef fishes (Gobiidae: *Eviota*). *Mol. Phylogenet. Evol.*, 82, 200–210. <https://doi.org/10.1016/j.ympev.2014.09.012>.
- Voris, H.K. (2000) Maps of Pleistocene sea levels in Southeast Asia: shorelines, river systems and time durations. *J. Biogeogr.*, 27, 1153–1167. <https://doi.org/10.1046/j.1365-2699.2000.00489.x>.
- Ward, R.D., Zemlak, T.S., Innes, B.H., Last, P.R. & Hebert, P.D.N. (2005) DNA barcoding Australia's fish species. *Philos. T. Roy. Soc. B*, 1462, 1847–1857. <https://doi.org/10.1098/rstb.2005.1716>.
- Whitfield, A.K., Panfili, A.K. & Durand, J.D. (2012) A global review of the cosmopolitan flathead mullet *Mugil cephalus* Linnaeus 1758 (Teleostei: Mugilidae), with emphasis on the biology, genetics, ecology and fisheries aspects of this apparent species complex. *Rev. Fish Biol. Fisher.*, 22, 641–681. <https://doi.org/10.1007/s1160-012-9263-9>.
- Xia, R., Durand, J.D. & Fu, C. (2016) Multilocus resolution of Mugilidae phylogeny (Teleostei: Mugiliformes): implications for the family's taxonomy. *Mol. Phylogenet. Evol.*, 96, 161–177. <https://doi.org/10.1016/j.ympev.2015.12.010>.
- Zemlak, T.S., Ward, R.D., Connell, A.D., Holmes, B.H. & Hebert, P.D.N. (2009) DNA barcoding reveals overlooked marine fishes. *Mol. Ecol. Resour.*, 9, 237–242. <https://doi.org/10.1111/j.1755-0998.2009.02649.x>.
- Zhang, J., Kapli, P., Pavlidis, P. & Stamatakis, A. (2013) A general species delimitation method with applications to phylogenetic placements. *Bioinformatics*, 29, 2869–2876. <https://doi.org/10.1093/bioinformatics/btt499>.