

Citrus Jabara Juice Diminishes Allergic Airway Inflammation in an Ovalbumin (OVA)-Induced Murine Asthma Model

Yasushi Ohno^{*}, BuLin BaiLa, Norihiko Funaguchi, Hirotoshi Iihara, Toshiya Inui, Yuka Sasaki, Daisuke Kawai, Chizuru Sakai, Daizou Kaitou, Koumei Yanase, Junki Endo, Shinya Mintoguchi and Hiroyuki Okura

Respirology of Internal Medicine Gifu University Graduate School of Medicine 1-1 Yanagido, Gifu 501-1194, Japan.

^{*}**Corresponding Author:** Yasushi Ohno, Respirology of Internal Medicine Gifu University Graduate School of Medicine 1-1 Yanagido, Gifu 501-1194, Japan. Tel: 81-58-230-6523, 81-58-230-6524, E-mail: yasusi@gifu-u.ac.jp

Citation: : Yasushi Ohno, BuLin BaiLa, Norihiko Funaguchi, Hirotoshi Iihara, Toshiya Inui, et al. (2023) Citrus Jabara Juice Diminishes Allergic Airway Inflammation in An Ovalbumin (OVA)-Induced Murine Asthma Model. *J Respir Dis Treat* 4(1): 102

Abstract

Background: Citrus jabara is a type of citrus fruit originally grown in Kitayama-Mura in Wakayama, Japan, and is reputed to have anti-allergic properties. In this study, we tested whether citrus jabara juice inhibits inflammatory mediators in a mouse allergic asthma model.

Methods: Mice were sensitized and challenged with ovalbumin to induce chronic airway inflammation. Two groups were orally administered citrus jabara juice at a daily dose of 10 mg/ml or 100 mg/ml in sterile water from day -3 to day 28.

Results: Administration of citrus jabara juice significantly reduced the numbers of infiltrating inflammatory cells in bronchoalveolar lavage fluid, as compared to control mice. Histopathological studies of the lung using hematoxylin & eosin and periodic acid-Schiff staining showed that citrus Jabara juice inhibited inflammatory cell infiltration and mucus hypersecretion in ovalbumin-challenged mice. In addition, the juice reduced levels of IL-4 while increasing levels of INF- γ .

Conclusions: These findings provide new insight into the immunopharmacological effects of citrus jabara and suggest its potential in supportive therapies for asthma.

Keywords: Citrus jabara juice; fruit; anti-allergy effects; bronchial asthma

Introduction

Flavonoids are polyphenolic plant secondary metabolites ubiquitously present in vegetables, fruits and beverages, and possess anti-allergic activities as well as immunomodulatory traits [1-9]. For example, flavonoids inhibit activation of mast cells and basophils, thereby suppressing the release of chemical mediators and the synthesis of Th2 type cytokines such as interleukin (IL)-4 and IL-13 [10,11].

Bronchial asthma is a complex disease of the lung characterized by reversible airway obstruction, allergic airway inflammation, excessive mucus production and airway hyper-responsiveness (AHR) [12]. Citrus jabara is a type of citrus fruit originally produced in Kitayama-Mura in Wakayama, Japan (Figure 1). Anecdotal evidence suggests citrus jabara possesses anti-allergic properties, but there is no scientific evidence at the moment. To address that issue, in the present study, we examined the degree to which citrus jabara juice protects against the development of ovalbumin (OVA)-induced bronchial asthma in Balb/c mice.

Fig. 1

Jabara



Figure 1 Jabara is a type of orange produced in Kitayama-Mura in Wakayama Prefecture, Japan

Methods

Animals and Experimental Protocol

Balb/c mice (6-8 weeks old) were purchased from Japan SLC (Sizuoka, Japan) and housed in separate cages according to the treatment protocol. Food and water were provided ad libitum. All experiments were carried out following the guidelines for the care and use of experimental animals formulated by the Japanese Association for Laboratory Animal Science in 1987.

Figure 2 illustrates the experimental protocol. Mice were sensitized to OVA by immunization on days 0 and 7 through intraperitoneal injection of 100 μ g of OVA (Grade V, Sigma Chemical Co. St. Louis, MO, USA) plus 1.6 mg of aluminum hydroxide in 200 μ l of 0.9% sterile saline (Otsuka, Tokyo, Japan). As a control, mice received only the aluminum hydroxide in saline. The OVA-sensitized mice were randomized into three groups (n=8 in each). Two groups were orally administered citrus jabara juice (Wakayama, Japan) at a daily dose of 10 mg/ml (Jabara L) or 100 mg/ml (Jabara H) in sterile water (0.3 ml/mouse) from day -3 to day 28. The third group received only the sterile water. The non-sensitized control mice (Sham; n=8) were also treated with sterile water only. The OVA-sensitized mice were then exposed to an aerosol of OVA (10 mg/ml) in 0.9% saline for 30 min each day on days 25, 26 and 27, whereas the non-sensitized mice were exposed only to saline. All mice were sacrificed 24 h after the last aerosol challenge.




Figure 2 Experimental protocol

Cell Counts in BALF

Bronchoalveolar lavage was performed by instilling 0.5 ml of saline into the lungs through a tracheal cannula and then gently aspirating the fluid. This was repeated three times. After staining with Wright's stain, total cell and eosinophil counts in the BALF were determined.

Peripheral Blood Cell Counts

Peripheral blood cell counts were made using blood samples collected from the orbital artery after the mice were sacrificed on day 28.

Histological Evaluation and Immunohistochemical Staining

After sacrificing the mice under anesthesia by intravenous injection with 30 mg/kg sodium pentobarbital on day 24, the lung tissues were fixed in 10% buffered formalin, embedded in paraffin, cut into 4-mm-thick sections and used for histological evaluation (H&E, PAS) or immunohistochemical staining. For immunohistochemical staining of IL-4, INF- γ and nitrotyrosine, phosphate-buffered saline (PBS) containing 0.05% Tween 20 and 2% normal goat serum was used as the antibody diluent after blocking endogenous peroxidase by addition of 0.3% H_2O_2 in methanol. The sections were incubated overnight at 4°C with human monoclonal antibodies against IL-4, INF- γ or nitrotyrosine (500:1 dilution; Oncogene Science, Cambridge, MA), washed with PBS to remove excess primary antibody, and then incubated with biotinylated horse anti-mouse IgG (200:1 dilution; Vector Laboratories, Burlingame, CA) for 1 h at room temperature. Bound antibody was then visualized using the standard protocol for the avidin-biotin-peroxidase complex protocol.

The degree of inflammation evident in the H&E-stained sections was scored as follows: 0, no sign of inflammation; 1, light or dispersed infiltrate in only a few areas of the section; 2, moderate infiltrate surrounding <50% of the vessels and airways; 3, heavy and focused infiltrate surrounding the majority of vessels and airways (13). Scoring for mucus production in PAS-stained sections was as follows: 0, no sign of mucus or increased numbers of PAS-positive cells in the airways; 1, no mucus in the airways, but a slight increase in the numbers of PAS-positive cells in a few airways; 2, some mucus is detectable in the airways, and ~50% of airway epithelial cells in multiple airways are PAS-positive; 3, several airways are plugged with mucus, and the majority of airway epithelial cells in multiple airways are PAS-positive. Scoring was carried out by two individuals blinded to the experimental

protocol.

Statistical Analysis

The data are presented as means \pm SD. Multiple comparisons among groups were made using one-way ANOVA with post hoc Tukey's tests. Values of $p < 0.05$ were considered significant.

Results

Cell Counts in BALF

We found that eosinophil counts were significantly higher in bronchoalveolar lavage fluid (BALF) from OVA-challenged mice than from saline-challenged control mice (Figure 3, A vs. B group). Moreover, administration of citrus jabara at a dose of 10 or 100 mg/ml (0.3 ml/day) significantly reduced eosinophil counts in BALF (Figure 3) from the OVA-challenged mice.

Fig. 3




Figure 3 Total cell and eosinophil counts in bronchoalveolar lavage fluid (BALF)

Lung Histology

To assess the histological effect of citrus jabara juice on allergen-induced airway inflammation, we analyzed lung tissues on day 24. In untreated, OVA-sensitized/challenged mice, numerous mucus-producing cells were observed within the airways, which were not seen in mice exposed only to aerosol saline. There was a substantial reduction in airway infiltration by PAS-positive cells in mice administered in citrus jabara juice at 100 mg/ml (0.3 ml/mouse), but not in those receiving citrus jabara at 10 mg/ml (0.3 ml/mouse) (Figure 4).

Fig. 4




Figure 4 Histological analysis of lung tissue sections stained with PAS

Immunohistochemical Staining

Numbers of periodic acid-Schiff (PAS)-positive mucus-producing cells within the airways were markedly higher in OVA-challenged mice than in control mice. Semiquantitative histological scores indicative of the degree of inflammation within lung tissue sections from OVA-challenged mice were significantly higher than the scores for corresponding control sections. OVA-induced infiltration of the lung tissue by inflammatory cells was significantly attenuated by 10 mg/ml or 100 mg/ml citras jabara juice, and there was a concomitant reduction in the inflammation scores in PAS-stained sections (Figure 5).

Histological Scores of PAS stain

Fig. 5




Figure 5 Histological scores for lung tissues stained with PAS

Histological analysis showed that airway infiltration by IL-4- and nitrotyrosine-positive cells was markedly greater in OVA-challenged than control mice, and that citrus jabara dose-dependently reduced staining for both IL-4 and nitrotyrosine in the lungs (Figure 6).




Figure 6 Histological analysis of lung tissues stained for IL-4

In addition, numbers of INF- γ -positive cells within the airways were markedly lower in the OVA-challenged mice than in the control mice, but citrus jabara juice reversed the effect of OVA on INF- γ expression within the lungs (Figure 7).




Figure 7 Histological analysis of lung tissues stained for INF- γ

Discussion

Bronchial asthma is characterized by acute and chronic airway inflammation and by AHR, the severity of which correlates with the degree of inflammation. In OVA-sensitized/challenged mice, repeated inhalation of antigen leads to increases in the levels of Th2 cytokines, including IL-4 and IL-5, and in the numbers of inflammatory cells (mainly eosinophils) in the BALF [14]. The cytokines expressed are involved in stimulating B cells to produce IgE and in promoting the infiltration of target tissues by mast cells and eosinophils [15]. IgE receptor-mediated activation of mast cells within the airways leads to oxygenation of arachidonic acid by 5-lipoxygenase and generation of leukotrienes (LTs), which promote airway edema, bronchial smooth muscle contraction and proliferation, enhanced mucus secretion and further eosinophil recruitment [16]. Several studies have shown that administration of TH1 cytokines, such as IFN- γ , at the time of sensitization can inhibit the induction of AHR and TH2-driven inflammation [17-19].

In humans, as in our mouse model, induction of IL-4 results in AHR, and blockade of IL-4 can relieve some of the symptoms of asthma [20]. IL-4 is central to the allergic response, promoting isotype switching of B cells to IgE synthesis, directing T cells along the Th2 differentiation pathway, upregulating the expression of vascular cell adhesion molecule-1 (VCAM-1), and controlling the expression levels of IgE Fc ϵ and the various cytokines and chemokine receptors involved in the allergic cascade. IL-4 drives IgE synthesis by B cells, and IgE levels in serum are dependent upon IL-4 [21]. In addition, IL-5 reportedly regulates the differentiation, recruitment and activation of eosinophils, which contribute to airway inflammation [22].

We previously reported that Narirutin is a flavonoid present in citrus jabara that exerts strong anti-allergic effects in a mouse model of asthma, and that the symptoms of allergic rhinitis are relieved by ingestion of the fruit juice ingredient of citrus jabara juice [23]. In the present study, we observed that citrus jabara significantly reduced allergen-induced airway inflammation in a mouse model of asthma, and that this effect was associated with significantly reduced IL-4 levels and increased IFN- γ levels within the lung tissues. We also found that citrus jabara juice reduced eosinophil counts in BALF, which could account for the reduced IL-4 levels. These studies show that administration of citrus jabara juice significantly diminishes such asthmatic reactions as leukocyte recruitment to the lungs, AHR and the production of Th2 cytokines, and that it attenuates inflammatory cell infiltration of lung tissue and goblet cell hyperplasia within airways. Taken together, these findings suggest citrus jabara may relieve airway inflammation, AHR and goblet cell hyperplasia through actions affecting production of Th1 and Th2 cytokines.

In summary, citrus jabara inhibited allergen-induced airway inflammation in a mouse model of asthma. The mechanism of its anti-asthma effect can be attributed, at least in part, to its ability to reduce IL-4 levels and increase IFN- γ levels. Citrus jabara thus has the potential for use in supportive therapies for bronchial asthma.

Acknowledgement

We thank Ms. Chieko Kashiwakura and Dr. Jun-ichiro Kan, Department of Orthopedic Surgery, and Dr. Daisuke Fujisawa, Department of Dermatology, Nihon University School of Medicine, Tokyo for their excellent technical assistance. We thank Drs. Kenko H. Lee and Masayuki Seki of Department of Orthopedic Surgery, Nihon University School of Medicine, Tokyo for their clinical sample collections. We thank Prof. Chisei Ra of Nihon University for his critical review and helpful suggestion. This work was supported in part by Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of the Japanese Government (Project No. [C] 15K09558, awarded to Y.O.), the Nihon University Multidisciplinary Research Grant for 2015 (Project No. So-16-014, awarded to Y.O.), and MEXT-Supported Program for the Strategic Research Foundation at Private Universities, 2015–2019 (Project No. S1511014, awarded to Y.O.).

The authors thank all participating sites with this study. This work was supported by Grant of Kitayama-mura, Wakayama, Japan.

Conflict of Interest

The authors have no conflict of interest

Author Contribution

YO, BB, NF and TG performed research and analyzed data. YO, YS and ST designed the study, DK, KY, FI, JE and HF performed the experiments. YO and SM wrote the paper.

References

1. Kimata M, Inagaki N, Nagai H (200). Effects of luteolin and other flavonoids on IgE-mediated allergic reactions. *Plant Med* 66: 25-9.
2. Higa S, Hirano T, Kotani M, Matsumoto M, Fujita A, et al. (2003) inhibits TH2-type cytokine production by activated human basophils. *J. Allergy Clin. Immunol* 111: 1299-306.
3. Hirano T, Higa S, Arimitsu J, Naka T, Shima Y, et al. (2004) Flavonoids such as luteolin, fisetin and apigenin are inhibitors of interleukin-4 and interleukin-13 production by activated human basophils. *Int. Arch. Allergy Immunol* 134: 135-40.
4. Williams CA, Grayer RJ (2004) Anthocyanins and other flavonoids. *Nat. Prod. Rep* 21: 539-73.
5. Chirumbolo S (2010) The role of quercetin, flavonols and flavones in modulating inflammatory cell function. *Inflamm. Allergy Drug Targets* 9: 263-85.
6. Visioli F, de la Lastra CA, Andres-Lacueva C, Aviram M, Calhau C, et al. (2011) Polyphenols and human health: A prospectus. *Crit. Rev. Food Sci. Nutr* 51: 524-46.
7. Calderon-Montano JM, Burgos-Moron E, Perez-Guerrero C, Lopez-Lazaro MA (2011) review on the dietary flavonoid kaempferol. *Mini Rev. Med. Chem* 11: 298-44.
8. Russo M, Spagnuolo C, Tedesco I, Bilotto S, Russo GL (2012) The flavonoid quercetin in disease prevention and therapy: Facts and fancies. *Biochem. Pharmacol.* 83: 6-15.
9. Magrone T, Jirillo E (2012) Influence of polyphenols on allergic immune reactions: Mechanisms of action. *Proc. Nutr. Soc.* 71: 316-21.
10. Kawai M, Hirano T, Higa S, Arimitsu J, Maruta M, et al. (2007) Flavonoids and related compounds as anti-allergic substances. *Allergol. Int.* 56: 113-23.
11. Matsuda H, Morikawa T, Ueda K, Managi H, Yoshikawa M (2002) Structural requirements of flavonoids for inhibition of antigen-induced degranulation, TNF-alpha and IL-4 production from RBL-2H3 cells. *Bioorg. Med. Chem.* 10: 123-6.
12. Busse WW, RF Lemanske, Asthma N (2001) *Engl. J. Med.* 344-50.
13. Kelly-Welch AE, Melo MEF, Smith E et al. (2004) Complex role of the IL-4 receptor α in a murine model of airway inflammation: Expression of the IL-4 receptor α on nonlymphoid cells of bone marrow origin contributes to severity of inflammation. *J. Immunol.* 172: 4545-55.
14. Paul MO, Byrne MB, FCCP (2006) Cytokines or their antagonists for the treatment of asthma. *Chest.* 130: 244-50.
15. Middleton EJ, Kandaswami C (1992) Effects of flavonoids on immune and inflammatory cell functions. *Biochem. Pharmacol.* 43: 1167-79.
16. Haczku A, K Takeda, I Redai, E Hamelmann, G Cieslewicz, et al. (1999) Anti-CD86 (B7.2) treatment abolishes allergic airway hyperresponsiveness in mice. *Am. J. Respir. Crit. Care Med.* 159:1638.

17. Gavett SH, O'Hearn DJ, Li X, Huang SK, Finkelman FD, et al. (1995) Interleukin 12 inhibits antigen-induced airway hyperresponsiveness, inflammation, and Th2 cytokine expression in mice. *J Exp Med.* 182: 1527-36.
18. Haapakoski R, Karisola P, Fyhrquist N, Savinko T, Wolff H, et al. (2011) Alenius H Intradermal cytosine-phosphate-guanosine treatment reduces lung inflammation but induces IFN- γ -mediated airway hyperreactivity in a murine model of natural rubber latex allergy. *Am J Respir Cell Mol Biol.* 44: 639-47.
19. Hirahara K, Yamashita M, Iwamura C, Shinoda K, Hasegawa A, et al. (2008) Repressor of GATA regulates TH2-driven allergic airway inflammation and airway hyperresponsiveness. *J Allergy Clin Immunol.* 122: 512-20.
20. Mark DA, CE Donovan, GT De Sanctis, SJ Krinzman, L Kobzik, et al. (1998) Both CD80 and CD86 co-stimulatory molecules regulate allergic pulmonary inflammation. *Int.Immunol* 10: 1647.
21. Finkelman FD, IM Katona, JF Urban, J Homes, J Ohara, et al. (1998) Interleukin 4 is required to generate and sustain in vivo IgE responses. *J. Immunol.* 141: 2335-41.
22. Humbert M, Corrigan CJ, Kimmitt P, Till SJ, Kay AB, et al. (1998). Relationship between IL-4 and IL-5 mRNA expression and disease severity in atopic asthma. *Am. J. Respir. Crit. Care Med.* 156: 704-8.
23. Funaguchi N, Ohno Y, La BL, Asai T, Yuhgetsu H, et al. (2007) Narirutin inhibits airway inflammation in an allergic mouse model. *Clin Exp Pharmacol Physiol.* 34: 766-70.

Submit your next manuscript to Annex Publishers and benefit from:

- ▶ Easy online submission process
- ▶ Rapid peer review process
- ▶ Online article availability soon after acceptance for Publication
- ▶ Open access: articles available free online
- ▶ More accessibility of the articles to the readers/researchers within the field
- ▶ Better discount on subsequent article submission

Submit your manuscript at

<http://www.annexpublishers.com/paper-submission.php>